

Original
ArticleLife
Science

Performance Characteristics of African Giant Land Snail (*Archachatina marginata*) Fed Diets Containing Soya Bean Milk Residue-Cassava Sievate Meal Mixtures

Olusegun Ojebiyi OJEBIYI^{1,*}, Omotayo Olugbenga JEJELOLA¹, Isaac Omoniyi OLADUNJOYE², Muritala Daniel SHITTU²

ABSTRACT [ENGLISH/ANGLAIS]

Thirty six growing African giant land snails (*Archachatina marginata*) with average initial live weight of 73.40±5.58 and 75.94± 4.70g were used in this study to evaluate the effect of diet containing soya bean milk residue (SBMR) and Cassava Sievate (CS) in ratio 3:2 on *Archachatina marginata*. The snails were randomly assigned to two treatments of eighteen snails each. The SBMR-CS mixture was included at 0 and 30% levels. Evaluated performance characteristics showed that daily feed intake, weight gain, feed-gain ratio, mortality, carcass analysis (foot weight, visceral weight, shell weight), shell characteristic (such as shell length, shell width), and mortality were not significantly ($p > 0.05$) affected by dietary treatments. The feed cost per weight gain (Nigerian Naira N/g) showed the low value of N 0.32 (0.002 USD) and high value of N 0.42 (0.003 USD) for the diets 2 and 1 respectively. It was concluded that SBMR/CS mixtures (3:2) can be included the diets of snails up to 30% without adverse effect on performance. The use of these agro by-products will support the ventures of increasing animal protein sources through cheaper snail production.

Keywords: Snails, soybean milk residue, cassava sievate, performance

RÉSUMÉ [FRANÇAIS/FRENCH]

Trente-six de plus en plus les escargots géants africains (*Archachatina marginata*) avec une moyenne de poids vif initial de 73,40 ± 5,58 et 75,94 ± 4,70g ont été utilisées dans cette étude pour évaluer l'effet de l'alimentation contenant des résidus de lait de soja haricot (SBMR) et le manioc Sievate (CS) 03h02 au ratio sur *Archachatina marginata*. Les escargots étaient assignés au hasard à deux traitements de dix-huit escargots chacun. Le mélange SBMR-CS a été inclus à 0 et les niveaux de 30%. Caractéristiques de performance évalués ont montré que l'ingestion quotidienne, gain de poids, l'alimentation du gain ratio, de mortalité, l'analyse des carcasses (poids du pied, le poids viscéral, poids de la coquille), coquille caractéristique (comme la longueur de la coquille, la largeur de coquille), et la mortalité n'étaient pas significativement ($p > 0,05$) affectées par les traitements diététiques. Le coût d'alimentation par le gain de poids (nairas nigériens N / g) a montré la faible valeur de N0.32 (0,002 \$) et la valeur élevée de N0.42 (américains Dollars 0,003 \$) pour les régimes 2 et 1 respectivement. On a conclu que SBMR / CS mélanges (03:02) peuvent être inclus les régimes alimentaires des escargots jusqu'à 30% sans effet négatif sur la performance. L'utilisation de ces sous-produits agricoles soutiendra le joint d'augmenter les sources de protéines animales grâce à une production moins chers d'escargot.

Mots-clés: Escargots, les résidus de lait de soja, le manioc sievate, la performance

Affiliations:

¹ Department of Animal Nutrition and Biotechnology, Ladoke Akintola University of Technology, PMB 4000, Ogbomoso, NIGERIA

² Department of Animal Production and Health, Ladoke Akintola University of Technology, PMB 4000, Ogbomoso, NIGERIA

* Email Address for Correspondence/ Adresse de courriel pour la correspondance: segunojebiyi@gmail.com

Accepted/Accepté: October, 2011

Full Citation: Ojebiyi OO, Jejelola OO, Oladunjoye IO, Shittu MD. Performance Characteristics of African Giant Land Snail (*Archachatina marginata*) Fed Diets Containing Soya Bean Milk Residue-Cassava Sievate Meal Mixtures. World Journal of Life Sciences and Medical Research 2011;1(5):105-10.

INTRODUCTION

Quantitatively and qualitatively, meat and other animal foods are better sources of proteins than plant foods (with the exception of soybean products) [1]. However, there is a great demand on the conventional animal protein food leading to their exorbitant prices and thus becoming unaffordable to some average Nigerians [2]. The desire to

increase animal protein intake therefore can be achieved through exploiting the potentials of non-conventional animal protein sources like snail. In terms of environmental friendliness, snail farming could be of choice. In addition, the low technical skill required for its rearing makes the venture acceptable to both urban and rural dwellers. The advocacy for snail rearing is becoming

more popular because of the potential usefulness of its meat as a good source of protein which is between 15.76 - 18.26 % [3, 4]. The fat content which ranged between 0.96 and 3.0% is low when compared with 9.6, 21.4 and 23.0 % in chicken, egg and mutton respectively [5]. This quality makes snail meat a choice for hypertensive patients and the obsessed [6, 7]. Snail meat is also very rich in Iron, magnesium, phosphorus, calcium and nitrogen [8]. The high content of iron in snail meat according to [9] makes it useful in correcting the problem of iron deficiency in man. However, as more people become aware of the medicinal and other importance of snail meat, snail might go into extinction with time [10] because the major source of snail supply to most Nigerian consumers is from the wild, which are sold along roadsides and in local markets. Hence, efforts are being made to domesticate snails intensively [11]. It has been reported by [12] that giant land snails fed plant food materials supplemented with artificial diet performed better than their counterparts fed only plant food materials. Feeding cost in livestock which accounts for over 70% of the total production costs can be reduced to ensure profitability through the use of agro by-products. Two of such by-products are Cassava Sievate (CS) and Soya Bean Milk Residue (SBMR) which are obtained from cassava and soya bean milk processing respectively. Their abundance, cheapness and availability all year round qualify them as good non-conventional ingredient [13].

This study is aimed at evaluating the effect of feeding diets containing graded levels of Soya Bean Milk Residue (SBMR)/Cassava Sievate (CS) meal mixture on the performance of growing snail.

MATERIALS AND METHODS

Location

The experiment was conducted at the Snailery Annex of Teaching and Research Farm of Ladoke Akintola University of Technology, Ogbomoso within the derived Savannah Zone of Nigeria. The agro-ecological description of the study area has been documented by Oguntoyinbo [14].

Preparation of Test Ingredients

Fresh cassava sievates were procured from the local Gari Processing Outlets within Ogbomoso and sundried for 5 days (Figure 1). Fresh Soya bean milk residues (SBMR) were procured from Soya milk processing outlets within Ogbomoso North Local Government Area. The residues

were immediately spread on clean slatted floor and sundried for 5 days under atmospheric temperature of 31 ± 2 °C (Figure 2).

The sundried SBMR and cassava sievate were hammer milled, bagged and stored until needed.

Preparation of Rearing Medium

Sandy loam soil collected from plots under cultivation within the Teaching and Research farm was sterilized and used as the rearing medium. Sterilization was done using hot water treatment. Water was heated to 100 °C and then poured into the medium (soil) to its level in plastic bowls used for the experiment. The water was later drained after cooling.

Formulation of the Experimental Diets

Two experimental diets were prepared. Diet₁ was the control without inclusion of CS-SBMR while Diet₂ had 30% of SBMR-CS mixtures in ratio 3:2. The diets were formulated to be isonitrogenous and isocaloric (Table 1).

Experimental Animal and Management

Thirty-six giant African land snail (*A. marginata*) growers were used for the experiment. The snails were fed with the control diet and acclimatized for one week before the commencement of the feeding trial. They were of average initial weight of 73.40 ± 5.58 - 76.02 ± 3.34 . Perforated plastic bowls of 20 cm depth and 40 cm diameter were used for housing the snails. They were filled with moist sterilized sandy loam soil to a depth of 15 cm, the basin were covered individually with mosquito net reinforced with chicken wire netting for proper aeration and to prevent escape of snails. They were kept under cool environmental condition.

After acclimatization, the snails were randomly allotted to two dietary treatments of eighteen snails per treatment. Each dietary treatment was replicated thrice at the rate of six snails per replicate in a completely randomized design experiment. The experiment lasted for 10 weeks, throughout which the snails were fed *ad-libitum* and were provided clean water in the evening (between 17-18 hour). While the experiment lasted, feed offered and orts were weighed to determine feed intake. Weekly weights were also taken after the initial weight and the records kept were used to determine the response criteria in term of mean weight gain, and feed intake to weight gain ratio. The shell length and shell width were also measured weekly using vener calipers.

Laboratory analysis

The proximate composition of the SBMR, CS and that of experimental diets were determined using the procedure of AOAC [15]. HCN determination was done using the procedure described by Bradbury et al. [16] and Egan et al. [17].

Statistical Analysis

All data generated were analyzed using Student T-test statistical analysis [18].

RESULTS AND DISCUSSION

The proximate composition of the test ingredients is presented in Table 2 while the proximate composition of the experimental diets is presented in Table 3. The proximate composition of the CS is higher than the value reported by Nwokoro et al. [19] but the crude fibre is lower. The experimental diets were similar in composition and the protein and energy values are similar to the values of Omole [20]. The protein is lower than that of Iman et al, Ueberu et al. [21, 22].

The effect of soya bean milk residue (SBMR)/Cassava Sievate (CS) meal mixture on performance characteristics of snail is presented in Table 4. The terminal weight, daily weight gain, daily feed intake were similar ($p > 0.05$) for snails on control diet and 30% inclusion of SBMR/CS mixtures respectively. There was no mortality in any treatment. The feed to gain ratio, feed cost per kg weight gain $\$0.003 \pm 0.00$ vs $\$0.02 \pm 0.00$ for snails on control diet and 30% inclusion of SBMR/CS mixtures respectively were affected ($p < 0.05$) by the dietary treatments.

The similarity in the terminal weights, daily feed intake, daily weight gain, and the zero mortality in both the treatment and control groups attest to the safety of the tested ingredients, as well as its potential usefulness in replacing maize. The final weights and daily weight gain obtained in this study were similar to the values reported by Oredein [23]. The zero mortality recorded in the snails fed control diet and 30% SBMR/CS mixture indicates that the inclusion of the test ingredients has no detrimental effect on the snails and confirms the report of Akinnusi [24] that, under proper management, the mortality rate in snails is lower than that of other conventional livestock.

The lower feed: gain ratio of diet 2 may be attributed to a better utilization of the diet. The shell morphological changes as presented in Table 4 shows that the terminal shell length, shell length increment, shell width and shell width increment were not affected ($p > 0.05$) by dietary treatments. This agrees with the findings of Amao et al.

and Omole [12, 20]. This could be due to the relatively similar ash content of the experimental diets.

The results of the carcass analysis are presented in Table 5. Although the visceral/live weight (%) as well as the shell/live weight (%) of the snails on the control diet did not differ ($p > 0.05$) from those of the snails on SBMR/CS-based diets, the values were lower than those reported by

Figure 1: This figure shows a flow chart of production of cassava sievate





Figure 2: This figure shows flow chart of production of Soy bean milk Residue



Odunaiya [25]. The results indicated that the foot weight which is otherwise referred to as edible portion was not affected ($p > 0.05$) by dietary treatment. The dressing

percentages of the snails in this study are similar for the control and the diet containing SBMR/CS, but are higher

Table 1: This table shows gross composition of experimental diets

Ingredients	% Composition	
	T ₁ (Control)	T ₂ (SBMR/CS)
Maize	20	2.35
Groundnut cake	14	14
SBMR/CS	0	30
Corn bran	14	1.65
Rice husk	16	16
Brewery dry grain	16	16
Palm kernel cake	14.25	14.25
Fish meal	1.0	1.0
Bone meal	2.0	2.0
Oyster shell	2.5	2.5
Vitamin premix	0.25	0.25
Total	100	100
Calculated analysis		
ME (Kcal/kg)	2499.5	2492.1
CP%	17.68	18.28
Fat	5.41	4.4
Fibre	9.82	8.65
Cost/kg of feed (N)	42.41	38.42

Premix Composition per kg of diet: Vitamin A – 3, 200, 000IU, Vitamin D₃- 1,200IU, Vitamin E₃- 200IU, Vitamin K₃- 800 mg, Vitamine B₁- 400mg, Vitamin B₂- 1000mg, Vitamin B₁₂- 2000 mg, Niacin- 400mg, Selenite (Se)- 40mg, Manganese (Mn)- 32,000 mg, Pantothenic Acid- 2,000 mg, Folic Acid-200mg, Choline Chloride- 60,000 mg, Iron (Fe)- 8,000 mg, Copper (Cu)-3,200mg, Zinc (Zn)-2,000 mg, Cobalt (Co)-90 mg, Iodine (I)-800mg; SBMR/CS – Soybean Milk Residue– Cassava Sievate mixtures; ME – Metabolizable Energy; CP – Crude Protein

Table 2: This table shows proximate composition of the test ingredients and experimental diets

Nutrients (%)	D1	D2	CS	SBMR	CSBM
Dry matter	89.94	89.54	88.25	90.35	89.65
Crude protein	17.40	17.39	2.34	29.94	1.81
Crude fibre	9.65	9.89	1.21	3.79	2.41
Ash	7.03	6.95	1.86	3.16	1.97
Ether extract	3.58	3.69	0.26	2.49	0.57
Nitrogen free extract	51.94	51.62	82.58	50.97	82.89
Gross energy (kcal/kg)	3.206	3.489	2.853	3.192	3.159
Cyanide content (mg/kg)	0.00	2.11	6.28	0.00	3.24

Cyanide content was determined using spectrophotometer method.

SBMR – Soya bean milk residue; D1 – Control diet; D2 – Diet with 30% inclusion level of SBMR/CS meal mixtures; CS – Cassava sievate; SBMR – Soybean milk residue; CSBM – Cassava sievate + soya bean milk residue (3:2).

Table 3: This table shows growth performance of snails fed diets containing graded levels of soya bean milk residue (SBMR) cassava sievate (CS) mixtures

Parameters	Treatments	
	1	2
Initial live weight (g)	73.40±5.58	75.94±4.70
Terminal live weight (g)	110.44±4.47	111.11±4.93
Average daily weight gain (g)	0.53±0.32	0.50±0.75
Total weight gain (g)	37.04±2.20	35.17±5.24
Feed/Gain ratio (FGR)	1.01±0.07 ^a	0.85±0.22 ^b
Mortality (%)	0.00	0.00
Feed cost (N/g)	0.042±0.00	0.038±0.00
¹ Feed cost/g weight gain(N)	0.42±0.00	0.032±0.22

^{ab}Means along the row with different superscript are significantly different (P<0.05); 1 – Control; 2 – 10% inclusion level of SBMR/CS meal mixtures; ¹One American dollar=One hundred and sixty Nigerian Naira

Table 4: This table shows Carcass analysis and shell characteristics of snails fed diets containing graded levels of soya bean milk residue (SBMR)/cassava sievate (CS) mixtures

Parameters	Treatments	
	1	2
Carcass Characteristics		
Mean live weight (g)	110.44±4.47	111.11±4.93
Foot weight (g)	48.69±2.75	46.61±2.31
Dressing (%)	44.09±1.79	42.00±1.69
Visceral weight	17.28±0.78	16.00±0.99
Shell weight (g)	18.18±1.68	22.19±1.30
Shell/live weight (%)	17.03±1.22	20.21±1.08
Visceral/live weight (%)	15.79±0.84	14.40±0.57
Foot/live weight (%)	40.17±1.07	38.73±0.76
Shell Characteristics		
Initial shell length (mm)	79.01±2.10	79.69±1.32
Terminal shell length (mm)	89.47±0.94	87.61±1.97
Shell length increment (mm)	10.46±1.43	8.83±1.90
Initial shell width (mm)	45.11±0.69	47.00±0.81
Terminal shell width (mm)	50.82±0.53	51.04±0.79
Shell width increment (mm)	5.71±0.50	4.49±0.77

SBMR – Soya bean milk residue; CS – Cassava sievate;

1 – Control diet; 2 – 10% inclusion level of SBMR/CS meal mixtures

than the values reported by Odunaiya, Omole [25, 26]. This may be an indication that the test ingredients used in this study has potential of producing more edible portion.

CONCLUSION

The study revealed that SBMR/CS mixtures are potential feed resource for replacing maize in snail feed. This will lower production cost, increase the farmers' profit margin,

and stimulate more snail production, hopefully, making more animal protein available for consumption.

REFERENCES

- [1] Heinz G, Hautzinger P. Meat processing technology for small- to medium scale producers. FAO/RAP Publication 2007/20. p. 456.
- [2] Ojelade AYP, Ibiyemi EA, Lamidi AW, Umoru JIA. Evaluation of performance characteristics of edible African Land snails (*Archachatina marginata*) fed different crude protein concentrate diets. In: Proceedings of the Nigerian society for Animal Production. 14-17 March, University of Ibadan, Nigeria. 2010. p. 236-9.
- [3] Awesu MO. The biology and management of Africa Giant Land snail (*Archachatina marginata*). M.Phil. Thesis, University of Ibadan. 1980.
- [4] Omole AJ. Utilization of Different Energy Supplement on the Performance Characteristics of Grower Edible Giant Land Snail (*A. marginata*). M.Sc. Dissertation, Anim. Sci. Department. Univ. of Ibadan. 1998.
- [5] Imevbore EA, Ademosun AA. The nutritive value of the African giant Land snail (*Archachatina marginata*). Journal of Animal Production Research 1998 (8) 2: 76-87.
- [6] FAO. Farming snails by FAO better farming series. 3/33 Rome Italy. 1986.
- [7] Odunaiya O. The utilization of snails in Nigeria. M.Sc. Thesis. University of Ibadan. 1995.
- [8] Hamzat RA, Omole AJ, Babatunde BB, Adejumo MO, Loge OG. Comparative value of kola testa, corn bran and rice bran with maize for mature African giant land snail (*Archachatina marginata*). Nigeria Journal of Animal Production 2007;34(2):265-76.
- [9] Cobbinah JR. Snail Farming in West Africa. The University of Chicago Press. C.T.A. Production (ACP EGG). London Convention. 1992.
- [10] Adeleke MA, Adeleye RB, Osinowo OA. Effect of Parent Body Weight on the Growth Rate of Giant African Land Snails Under Unrestricted Feeding. In: Proc. of 7th Ann. Conf. of Anim. Sci. Asso. of Nigeria (ASAN). September, 16-19, Uni. of Agric. Abeokuta (UNAAB), Nigeria. 2002. p. 369-72.
- [11] Kehinde AS, Adebayo O, Usman JM, Meduna AJ. Effect of Incorporation of Cassava papaya Leaf and Fruit in Ration of Snails on the Growth Performance Characteristics Proc. of 27th Ann. Conf. of Nigeria Society for Ann. Pro. (NSAP). March 17-21, Federal University of Technology, Akure, Nigeria. 2002. p. 319-20.
- [12] Amao OA, Oladunjoye TO, Aderogba OA. Effect of Supplementing Fresh Pawpaw Leaf Diet with Different Milling By-products on the Growth Performance of African Giant Snail (*Archachatina marginata*). In: Proceedings of 7th Annual Conference, Animal Science Association of Nigeria (ASAN). September 16-19, University of Agriculture, Abeokuta (UNAB), Nigeria. 2002. p. 277-80.
- [13] Ojebiyi OO, Farinu GO, Babatunde GM. Nutritional Potentials of Differently Processed cassava peels and bloodmeal as livestock feed ingredients. Journal of Agriculture, Forestry and the Social Sciences 2010;8(1):195-207.
- [14] Oguntoyinbo JS, Climate. In: Oguntoyinbo JS et al (eds). Geography of Nigeria Development, Heinemann Nigeria Ltd, Ibadan; 1988. p.78.
- [15] AOAC. Association of Official Analytical Chemists. Official Methods of Analysis. 15th (Edn.), Washington D. C. USA, 1990. p. 69-88.
- [16] Bradbury MG, Egan SV, Bradbury JH. Determination of all forms of cyanogens in cassava roots and cassava products using picrate paper kits. Journal of Science of Food and Agriculture 1999;79:593-601.
- [17] Egan SV, Yeoh HH, Bradbury JH. The cyanogenic potential of cassava Journal of Science of Food and Agriculture 1988;76:39-48.
- [18] Steel RGD, Torrie JH. Principles and Procedures of Statistics. A biometrical Approach. 2nd Edition. MacGraw Hill. Kogakasha Limited. 1990.
- [19] Nwokoro SO, Vaikosen SE, Bamgbose AM. Nutrient composition of cassava offals and cassava sievates collected from locations in Edo State, Nigeria. Pakistan Journal of Nutrition 2005 4(4): 262- 264.
- [20] Omole AJ. Nutritional evaluation of cassava Peel-palm kernel cake mixtures in snails diet. Proceedings of the 36th Annual conference of the Nigerian society for animal production 13-16th March, University of Abuja. 2011 p305-307.
- [21] Iman GT, Okunade SA, Adebago CFA. Effect of replacing soya and fish-meal with mulberry and sian-weed leaves in the supplementary diet of

- Archachatina marginata Swais. (African Giant Snail). In: Proceedings of the 14th Annual conference of ASAN 14th 16th September, Ladoko Akintola University of Technology, Nigeria. 2009. p. 567-8.
- [22] Uberu NP, Ani AO, Foleng-Ndofor HM, Egenti O. Performance of African Giant Snail (*Archachatina marginata*) fed varying levels of Bambara offal. Proceedings of the 35th Annual Conference of NSAP, 14-17 March, University of Ibadan, Nigeria. 2010. p. 260-2.
- [23] Oredein AO, Omole AJ, Ognidele FO. Performance characteristics of growing snails (*Archachatina marginata*) fed different Agro-by products. Proceedings of the 28th Annual conference of NSAP. 2003. p. 325-54.
- [24] Akinnusi O. A Practical Approach to Background Snail Farming. Nigeria Journal of Animal Production 1998;25:1-3.
- [25] Odunaiya O, Akinyemi AA. Performance of two snails species- (*Archachatina marginata*) and *Achatina achatina* reared under the same management practices. Nigerian Journal of Animal Production 2008;35(2):224-9.
- [26] Omole AJ. Performance and carcass analysis of growing snails fed *Calopogonium mucunoides* (Calopo) and *Pueraria phaseoloides* (Kudzupuro). Nigerian Journal of Animal Production 2010;37:85-90.

ACKNOWLEDGEMENT / SOURCE(S) OF SUPPORT

Nil.

CONFLICT OF INTEREST

No conflict of interests was declared by authors.

How to Submit Manuscripts

Since we use very fast review system, and since we are dedicated to publishing submitted articles with few weeks of submission, then the easiest and most reliable way of submitting a manuscript for publication in any of the journals from the publisher Research, Reviews and Publications (also known as Research | Reviews | Publications) is by sending an electronic copy of the well formatted manuscript as an email attachment to rrpjournals@gmail.com or uploaded at <http://rrpjournals.com/blog/SUBMIT-MANUSCRIPT.php>.

Submissions are often acknowledged within 6 to 24 hours of submission and the review process normally starts within few hours later, except in the rear cases where we are unable to find the appropriate reviewer on time.

Manuscripts are hardly rejected without first sending them for review, except in the cases where the manuscripts are poorly formatted and the author(s) have not followed the instructions for manuscript preparation which is available on the page of Instruction for Authors in website and can be accessed through <http://www.rrpjournals.com/InstructionsForAuthors.html>.

Research | Reviews | Publications and its journals have so many unique features such as rapid and quality publication of excellent articles, bilingual publication, some of which are available at <http://www.rrpjournals.com/uniqueness.html>.