

Original Article

Public Health

Microbial Contaminants of Wooden Toothpicks in Abakaliki Metropolis, Ebonyi State, Nigeria

Michael Okpara ELOM¹, Uchenna Iyioke UGAH², Victor OMOTE¹

ABSTRACT [ENGLISH/ANGLAIS]

Evaluation of microbial contaminants of wooden toothpicks was carried out in Abakaliki metropolis, of Ebonyi State, Nigeria. A total of one hundred (100) samples comprising of ten (10) toothpicks each from four different sources within six zones were analyzed for microbes, using wet preparation techniques, concentration techniques, culture, and biochemical tests. Out of the 80 (eighty) test samples analyzed, 53 (66%) showed microbial contamination while 27 (34%) were sterile. Of the 53 positive samples, 24 (45%) gave mixed contaminations while 29 (55%) were contaminated by single microbes. Bacteria accounted for the highest contamination (positive for 42 samples out of the 53) while parasites were recovered from 2 samples only. Ogoja Road and Ishieke village showed the highest rates of contamination with 78% and 77% respectively while PRESCO and CAS campuses gave 57% and 54% contaminations respectively. Samples collected from restaurants showed 86% contamination while those collected from eateries gave 54% contamination. Yeast cells and fungal spores contaminated 22 (twenty-two) and 18 (eighteen) of the 53 positive samples respectively while staphylococcus aureus, Escherichia coli and Pseudomonas aeruginosa were the most recorded bacteria with prevalence of 8 (eight) each. Proteus and Salmonella contaminated 2 (two) samples each while Giardia lamblia and Ascaris lumbricoides were isolated from one sample.

Keywords: Toothpick, contaminants, restaurants

RÉSUMÉ [FRANÇAIS/FRENCH]

Évaluation des contaminants microbiens de cure-dents en bois a été réalisée en métropole Abakaliki, de l'Etat d'Ebonyi, Nigeria. Un total de cent (100) des échantillons comprenant des dix (10) des cure-dents chaque de quatre sources différentes dans les six zones ont été analysés pour les microbes, en utilisant des techniques humides de préparation, les techniques de concentration, la culture et les tests biochimiques. Sur les 80 (quatre-vingts) échantillons d'essai analysés, 53 (66%) ont montré une contamination microbienne tandis que 27 (34%) étaient stériles. Sur les 53 échantillons positifs, 24 (45%) a donné des contaminations mixtes tandis que 29 (55%) ont été contaminés par des microbes simples. Bactéries représentaient la plus forte contamination (positif pour 42 échantillons sur 53), tandis que les parasites ont été récupérés à partir de 2 échantillons seulement. Ogoja route et le village Ishieke ont montré des taux de contamination avec 78% et 77% respectivement, tandis que les plus élevés PRESCO et CAS campus ont donné 57% et 54% respectivement des contaminations. Les échantillons prélevés dans les restaurants ont montré la contamination de 86% tandis que celles recueillies auprès de restaurants a donné la contamination de 54%. Les cellules de levure et des spores fongiques contaminés 22 (vingt-deux) et 18 (dix-huit) des 53 échantillons positifs respectivement, tandis que le staphylocoque aureus, Escherichia coli et Pseudomonas aeruginosa étaient les bactéries les plus enregistrés avec une prévalence de 8 (huit) chacun. Proteus et Salmonella contaminés deux (2) échantillons chacun tandis que Giardia Lambia et Ascaris lumbricoides ont été isolés à partir d'un échantillon.

Mots-clés: Cure-dents, les contaminants, les restaurants

Affiliations:

¹ Department of Medical Laboratory Science, Faculty of Health Sciences and Technology Ebonyi State University, Abakaliki

² Department of Medical Biochemistry, Faculty of Basic Medical Sciences, Federal University, Ndufu-Alike Ikwo

* Email Address for Correspondence/
Adresse de courriel pour la correspondance:
ugahuchennaiyioku@gmail.com

Accepted/Accepté:
April, 2014

Full Citation: Elom MO, Ughah UI, Omote V. Microbial Contaminants of Wooden Toothpicks in Abakaliki Metropolis, Ebonyi State, Nigeria. World Journal of Life Science and Medical Research 2014;3(3):101-6.

INTRODUCTION

Microorganisms are ubiquitous in nature. They live in every part of the biosphere including soil, hot springs, on the ocean floor, in the atmosphere and deep inside rocks within the earth crust [1]. Microbes are very important tools that have been exploited by people in biotechnology, both in traditional food and beverages preparation, and in

modern technologies based on genetic engineering. However, there are many pathogenic microbes, which are harmful and can even cause death in plants and animals [2]. The pathogenic microbes possess various pathogenic mechanisms that enable them to gain access into the host, attach to receptors on cell surfaces, replicate locally,

escape host immune responses, spread to different parts of the body and cause cellular and tissue destructions.

A toothpick is a tiny piece of wood, plastic, bamboo, metal, bone and other substances used to remove debris from the teeth, usually after a meal. Toothpicks usually have one or two sharp ends to insert between teeth. They come in different shapes, sizes, colours, and may also have different flavors. Those made from metals, plastics and wood are the most common types found. Besides being used to remove food from and between teeth, they can be used to perform a number of functions ranging from designing building plans in architecture, to supporting broken stems in floriculture, cleaning and painting hard to reach or awkward places, cake baking and designing, cleaning of nails, applying glue to tight spots and others [3]

Toothpicks are one of the most poorly handled eating accessories. They are usually placed in bowls or plates and left open in restaurants, bars, eateries and other places where they are used. This unhealthy habit renders them exposed to contaminants from the air, hands, skin and vectors of microbes, such as the houseflies and rodents. The contaminated toothpicks serve as fomites, transmitting the microbes, which have the potentials of causing serious public health problems.

The fear that toothpicks could serve as germ spreaders because of how they are usually exposed in bowls or cups where restaurants patrons often recklessly handle them, fingering many and leaving them behind, started gaining attention in the early 20th century and led to their elimination from dinner tables and the ban of open containers of toothpicks by health authorities in Minneapolis [4]. The propounded inherent danger in the use of toothpicks gained support through the works of Chang and colleagues in which pathogenic bacteria were isolated from hands injured by toothpicks [5].

Recovery of microorganisms from toothpicks in restaurants, bars, eateries etc. signifies a looming public health danger, considering the nutritional, immunological and physiological effects of the microbes on their human hosts. There is little information on the health implications of toothpicks as fomites. This can be largely attributed to the fact that its use has reduced drastically over the years because most elite societies see its use as vulgar and as an offensive and uncivilized practice and most researchers often neglect the pathogenic-bearing potentials of toothpicks. The dearth of information on the pathogenic

transmission potentials of toothpicks encouraged the execution of this study, so as to close the information gap.

MATERIALS AND METHODS

Study Area

The study was carried out in Abakaliki metropolis in Ebonyi State, South –eastern Nigeria. Abakaliki is the capital of Ebonyi State, Nigeria. The climate is tropical with two distinct seasons (rainy and dry). The former commences from April and ends in October while the latter starts from November and ends in March. The average rainfall of the city is about 1500 mm while the temperature range is 28-30°C. The city is located very close to the surrounding rural communities where mostly farming activities take place.

Sampling Techniques

A total of one hundred (100) samples (each comprising of ten (10) toothpicks) from four (4) different sources were used in the study. The sources were restaurants, bars, eateries and the market. The study area was divided into six zones namely PRESCO campus and its environs, CAS campus and its environs, Water-works Road and its environs, Ogoja Road, Federal Teaching Hospital Abakaliki (FETHAI) and its environs and Ishieke village. Eighty (80) samples (800 toothpicks) out of the total population were test samples while the remaining twenty samples (200 toothpicks) were bought from the market and used as control samples. The samples were collected from July through September, 2013, using simple random sampling technique. Each of the collected toothpicks were placed in a sterile polythene bag (newly bought drug envelop) sealed and transported to the Research Laboratory in the Department of Medical Laboratory Science, Ebonyi State University, Abakaliki, for analysis.

Laboratory Analysis

The laboratory methods and techniques followed WHO [6]. The toothpicks were placed in a sterile universal container containing 10 ml of normal saline solution and the container was shaken vigorously. Thereafter, a sterile forcep was used to remove the toothpicks. The solution was poured into a centrifuge tube and centrifuged at 2,000 g for 5 minutes. The supernatant was discarded leaving behind the sediment. A part of the sediment from the centrifuged solution was used to aseptically inoculate blood agar and MacConkey agar plates as described by

Barker and Silverton [7] and was incubated for 24 hours at 37°C.

A small quantity of the sediment was placed on a clean, dry and grease-free glass slide. The preparation was covered with a cover glass and examined under the microscope using x10 and x40 objectives respectively.

After 24 hours of incubation, the cultured plates were examined macroscopically for growth of any microbes. The microbial isolates were identified and characterized using standard methods [8].

Statistical Analysis

The analysis of recovered and isolated microbes was carried out using descriptive statistics and inferential statistics of Chi-Square.

RESULTS

Out of the 80 test samples, 53 (66%) showed microbial contamination while 27 (34%) were sterile. The 20 control samples showed no microbial contamination of any sort.

Table 1: Prevalence of contamination in relation to zones

Zones	Sample size (n)	Number of contaminated samples	Percentage of contaminated samples
Ogoja Road	14	11	78.57
Ishieke Village	13	10	76.92
Water-works Road	13	9	69.23
FETHA1 and Its environs	13	8	61.54
PRESCO Campus	14	8	57.14
CAS campus	13	7	53.85

FETHA 1: Federal Teaching Hospital, Abakaliki.

PRESCO campus: College of Health Sciences, Ebonyi State University

CAS: College of Agricultural Sciences

Forty-five percent (45%) of the contaminated samples gave mixed contamination (contained more than one microbe) while 55% were contaminated by only one microbe. Bacterial contamination was found in 42 of the 53 contaminated samples (79%). Fungi contamination was

found in 33 samples (63%) while only 2 samples (3%) were contaminated by parasites. Ogoja Road had the highest contamination while CAS campus had the least (Table 1). Of the three sources of sample, restaurants had the highest contamination while eateries had the least contamination (Table 2).

The prevalence of contamination with respect to microbial species is presented in table 3. Yeast cells had the highest prevalence while the parasites (*Giardia lamblia* and *Ascaris lumbricoides*) had the least.

Table 2: Microbial contamination in relation to sources

Sources	Sample size	Number of Contaminated samples	Percentages of Contaminated samples
Restaurants	29	25	86.21
Bar	26	15	57.69
Eatery	25	13	52.00

Table 3: Prevalence of contaminants with respect to microbial species

Microbes	Number of contaminated samples	Percentage of contaminated samples
Yeast cells	22	41.51
Fungal spores	18	33.96
<i>Staphylococcus aureus</i>	8	15.09
<i>Escherichia coli</i>	8	15.09
<i>Pseudomonas aeruginosa</i>	8	15.09
<i>Non-coagulase Staphylococci</i>	5	9.43
<i>Streptococci spp</i>	5	9.43
<i>Klebsiella spp</i>	4	7.55
<i>Salmonella spp</i>	2	3.77
<i>Proteus spp</i>	2	3.77
<i>Giardia lamblia cyst</i>	1	1.89
<i>Ascaris lumbricoides</i>	1	1.89

DISCUSSION

Out of the 80 test samples, 53 (66%) showed contamination. This high prevalence of contamination could be attributed to the fact that the toothpicks are made from woods which are naturally porous and have tiny fissures and grooves that can harbor microbes, particularly bacteria [9]. This fact was supported by the

work of Gilbert and Watson [10] which showed that microbes were more abundant on wooden boards when compared to metal or plastic surfaces after contact with food products. The fact that the study area (Abakaliki) is surrounded by rural communities [11] where sanitation and personal hygiene is low might also be a contributing factor to the high prevalence of the microbes recorded. Forty-five percent (45 %) of the contaminated samples showed mixed contamination. This work agrees with the findings of Lam and Choi [12] in which more than one bacterium species were isolated from the sample of patients injured by toothpicks.

Bacteria had the highest level of contamination, followed by fungi, while the parasites had least. A number of factors might be responsible for this but of paramount importance might be the species of woods involved. Bacteria recovered from birch wood after contamination with *E.coli* for 3 days, indicated that birch wood has reduced antibacterial effect when compared to other species of wood [13]. Moisture content, temperature and humidity are the key factors for fungal growth on different materials [9]. This may account for the high prevalence of fungi and low prevalence of parasites recovered in the study, as Abakaliki is known for such environmental factors.

Ogoja Road and Ishieke village gave the highest rate of contamination of 78% and 77% respectively while PRESCO and CAS campuses showed contaminations of 57% and 54% respectively. Ogoja Road is the heart of commercial activities in Abakaliki metropolis and the home to most small scale business firms in Ebonyi State. The increased commercial activities as well as the low standard of infrastructure in this area help to promote poor sanitation and poor personal hygiene which may be responsible for the high prevalence of microbial contaminations observed. Ishieke village is inhabited by most natives and few students of the Faculties of Education, Management Sciences, Social Sciences, and Arts and Humanities of Ebonyi State University, Abakaliki. Ignorance, unwholesome practices and lack of basic social amenities in the area might be responsible for poor sanitary conditions which may have accounted for the high prevalence of toothpick contaminations recorded. The reduced prevalence observed in both PRESCO and CAS campuses may be attributed to health education and enlightenment which are factors that promote personal hygiene and improved sanitation. Most persons residing in those areas were students, lecturers, some members of

university staff, a few traders and other individuals with educational exposures and enlightenments.

With respect to sources, the highest rate of microbial contamination was observed in restaurants. This might be due to the fact that most restaurants in Abakaliki are below standard. The restaurants are managed by people with poor personal hygiene and low sanitary values as well as being patronized by individuals with poor knowledge, attitude and practices of unwholesome habits. Most bars in the study area are combined with restaurants. This might be responsible for the prevalence of 58% contamination recorded for bars.

Yeast cells and fungal spores were the most recorded microbes. This work agrees with Valentine [14] who showed that fungal species contaminated Archives and Museums than bacteria species. This may be attributed to the strong cellulolytic activity exerted by fungi on wood [15]. *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa* showed prevalence of 15% contamination each. This study is in consonance with that of Famurena and David [16] which revealed that the above-named bacteria showed the highest prevalence among bacteria that were recovered from cell phones. The high prevalence of *Staphylococcus aureus* may be attributed to the fact that it can remain viable for about 9 weeks on dry surfaces [17]. It is also estimated that 20 % of the human population are long-term carriers of *S. aureus* and that it is a normal flora of the skin [18]. This might also be a contributing factor to the high prevalence recorded. Fecal-oral transmission is the major route through which strains of *E. coli* are transmitted [19]. Thus, poor personal hygiene such as non-washing of hands after passing out faeces and fecal contamination of water sources may be responsible for the high prevalence of *E.coli* in the study.

P. aeruginosa is a ubiquitous organism present in diverse environmental settings [20]. Its ability to survive on minimal nutritional requirements and to tolerate a variety of physical conditions [20] may be responsible for the high prevalence of *P. aeruginosa* on toothpicks. The survival of parasite (cysts, ova, trophozoites or adult worms) in the environment depends on both biotic and abiotic factors [21]. The biotic factors include other microbes and their by-products while the abiotic factors include temperature, relative humidity and residence time [21]. *E. histolytica* can survive for only 18 hours on dry surface where as it can survive for 72 hours and up to 10 days on moist and damp surfaces respectively [22]. The poor contamination of the studied toothpicks by parasites may be as a result of

increased contamination by other microbes and the dry nature of the toothpicks studied. The only helminth recovered in the study was *Ascaris lumbricoides*. This may be attributed to the fact that *Ascaris* eggs are more resistant than other intestinal parasite eggs [21] and the mucopolysaccharide that coats the eggs serves as an adhesin that aids its attachment to surfaces [23].

Statistical analysis using Chi-square at a significant level of 95% showed that the prevalence of the recovered microbes is independent of the studied zones and the sources of the sample. The high prevalence of microbial contaminants recorded in the study, indicates a looming public health danger. Most of the microbes recovered were pathogens that could be very destructive on their hosts and are also implicated in the growing antimicrobial resistance phenomenon that is posing a threat to chemoprophylaxis and chemotherapeutics of infectious diseases. Recovered microbes such as yeast and some fungal species that had not been termed pathogens in the past are now threats to the health of immunocompromised individuals. Further research that may lay emphasis on the quantitative analysis of the microbes is recommended. Such a study may find out the least quantities of microbial counts that could establish asymptomatic pathogenic effects on users of toothpicks.

Poor sanitation practices and personal hygiene have been incriminated as the main factors responsible for the high prevalence of microbial contaminants in this study. Therefore, government and relevant authorities should ensure the provision of basic social amenities, as well as the enactment and enforcement of health laws that would reduce microbial contamination of objects and surfaces in restaurants, bars and eateries. Health education and enlightenment programmes on preventive and control measures of infectious diseases and prompt diagnosis and treatment of infected persons should also be emphasized. Regular seminars should always be organized for restaurant managers and other food sellers. Toothpicks made of other materials or other wood species with potent antimicrobial activities are recommended in preference to the common ones made from birch wood.

REFERENCES

- [1] Ronniie G, Frank W, Mathias M, Kazumasa O, Robert T, Donald EC, Hirishi K. High rates of microbial carbon turn over in sediments in the deepest oceanic trench on earth. *Nature Geoscience* 2013;6(4):284.
- [2] Talaro PK, Chess B. *Foundation in Microbiology*, 8th edition. Mc-Graw Hill Company, Incorporated, New York. 2008. Pp 3-8.
- [3] Petroski H. *The Toothpick: Technology and Culture*. Knop Double Day Publishing Group. New York. 2007. Pp 6-46.
- [4] Whitaker J. *Toothpicks. Restauranting through history*. American Palate 2012;13:27-32.
- [5] Chang CM., Huang LY, Liu HP, Lo HW. Infectious complications associated with toothpicks injuries of the hand. *Journal of Hand Surgery* 2003;28(2):327-31.
- [6] World Health Organization. *Manual of Basic Technologies for a Health Laboratory*, 2nd edition, Geneva, Switzerland. 2003.
- [7] Baker FJ, Silverton RE. *Introduction to Medical Laboratory Technology*, 7th edition. Bounty Press Limited, London. 1998. Pp. 276.
- [8] Chessbrough M. *District Laboratory Practice in Tropical Countries*, part 1. Cambridge University Press, London. 2000. Pp. 193.
- [9] Annett M., Rolf K, Alfred, W, Kornella S. Survival of Bacteria on Wood and Plastic Particles: Dependence on wood species and environmental condition. *Halzforschung* 2005;59:72-81.
- [10] Gilbert R, Watson JHM. (1971). Some laboratory experiments on various meat preparation surfaces with regards to surface contamination and cleaning. *Journal of Food Technology*, 6: 163-170.
- [11] Elom MO, Alo MN, Ezike AC, Okeh EN, Anyim C. Parasitic helminthes on Nigerian currency: a public health jeopardy. *Prime Journal of Microbiology Research* 2012;2(6):165-9.
- [12] Lam, H. and Choi, Y. (2004). Pathogenic hand infections: importance of Gram-negative organisms. *Hong kong Journal of Orthopedic Surgery*, 8 (1) : 28-32.
- [13] Ak NO, Clivor DO, Kasper CW. Cutting boards of plastic and wood contaminated experimentally with bacteria. *Journal of Food Protocol* 1994;57:16-22.
- [14] Valentine M. *Microbial contamination in Archives and Museums: Health Hazards and Preventive Strategies using Air Ventilation Systems*. Contribution to the Experts Round-table on Sustainable Climate Management Strategies. Temerise, Spain. 2007.

- [15] Anna N, Rrafai LG, Angenieszka W. Microbial contamination of Store.Rooms at the Aushwitz-BirkenauMuseum. *Aerobiologia* 2010;2:125-33.
- [16] Famurewa O, David OM. Cell phone: a medium of transmission of bacteria pathogens. *World Rural Observations* 2009;1(2):68-72.
- [17] Beard-Peglear MA, Stubbs E, Vickery AM. Observation on the resistance to drying of Staphylococcal Strains. *Journal of Medical Microbiology* 1988;26:251-5.
- [18] Kluytmans J, Van-Belkum A, Verbrugh H. Nasal Carriage of Staphylococcus aureus: Epidemiology, underlying Mechanisms and associated risks. *Clinical Microbiology Reviews* 1997;10(3):515-20.
- [19] Feng P, Weagent S, Grant M. Enumeration of Escherichia coli and the coliform bacteria. *Bacteriological Analytical Manual*. 8th edition. Center for Food Safety and Applied Nutrition. 2002.
- [20] Lister DP, Wolter JD, Hanson DN. Antibacterial resistant Pseudomonas aeruginosa: clinical impact and complex regulation of chromosomally-encoded resistance mechanisms. *Clinical Microbial Reviews*, 2009;22(4):582-610.
- [21] Alum A, Rubiono KJ, Ijaz KM. The global war against intestinal parasites- should we use a holistic approach? *International Journal of Infectious Diseases* 2010;14:732-8.
- [22] Feachem RG, Bradley DJ, Garelick A, Mara DD. *Appropriate Technology for Water Supply and Sanitation: health aspect of excreta and sewage management. A State-of- the Art Review, Volume 3.* Washington D C. The World Bank. 1981.
- [23] Crompton DWT. *Biology of Ascaris lumbricoides.* In: Crompton DWT, Neshemm MC, Pawlowsski ZS. (Editors). 2nd Edition. *Ascaris and its Prevention and Control.* Tailor and Francis Publisher, London. 1989. Pp 9-14.

ACKNOWLEDGEMENT / SOURCE(S) OF SUPPORT

Please include here any form of relevant acknowledgement including those to the contributors who are not authors.

CONFLICT OF INTEREST

Please kindly declare any conflict of interest here

How to Submit Manuscripts

Manuscript must be submitted online. The URL for manuscript submission is <http://rrpjournals.org/submit>

Manuscript submissions are often acknowledged within five to 10 minutes of submission by emailing manuscript ID to the corresponding author.

Review process normally starts within six to 24 hours of manuscript submission. Manuscripts are hardly rejected without first sending them for review, except in the cases where the manuscripts are poorly formatted and the author(s) have not followed the guidelines for manuscript preparation, <http://rrpjournals.org/guidelines>

Research | Reviews | Publications and its journals (<http://rrpjournals.org/journals>) have many unique features such as rapid and quality publication of excellent articles, bilingual publication, and so on.